



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## SOLUTIONS OF PROBLEMS.

**2669. Proposed by S. A. COREY, Albia, Iowa.**

Let  $A_1, A_2, \dots, A_8$ , and  $-(A_1 + A_2 + \dots + A_8)$  be the vector sides of an enneagon, plane or gauche. Also let  $B_1, B_2, \dots, B_8$ , and  $-(B_1 + B_2 + \dots + B_8)$  be the vector sides of a second enneagon, where  $B_1 = C_1A_1 - C_2C_6A_3 - C_3C_6A_5 + C_4C_6C_6A_7$ ,  $B_2 = C_1A_2 - C_2C_6A_4 - C_3C_6A_6 + C_4C_6C_6A_8$ ,  $B_3 = C_2A_1 + C_1A_3 - C_4C_6A_5 - C_3C_6A_7$ ,  $B_4 = C_2A_2 + C_1A_4 - C_4C_6A_6 - C_3C_6A_8$ ,  $B_5 = C_3A_1 + C_4C_6A_3 + C_1A_5 + C_2C_6A_7$ ,  $B_6 = C_3A_2 + C_4C_6A_4 + C_1A_6 + C_2C_6A_8$ ,  $B_7 = C_4A_1 - C_3A_3 + C_2A_5 - C_1A_7$ , and  $B_8 = C_4A_2 - C_3A_4 + C_2A_6 - C_1A_8$ ,  $C_1, C_2, C_3, C_4, C_5$ , and  $C_6$  being scalars.

Then, if  $a_s$  = tensor  $A_s$ ,  $b_s$  = tensor  $B_s$ , and  $\cos(A_rA_s)$  = cosine of the angle included between  $A_r$  and  $A_s$  and  $\cos(B_rB_s)$  = cosine of the angle included between  $B_r$  and  $B_s$ , establish the following relation between the sides and angles of the two enneagons:

$$\begin{aligned} [C_1^2 + C_5C_2^2 + C_6C_3^2 + C_5C_6C_4^2][a_1a_2 \cos(A_1A_2) + C_5a_3a_4 \cos(A_3A_4) + C_6a_5a_6 \cos(A_5A_6) \\ + C_5C_6a_7a_8 \cos(A_7A_8)] \\ = b_1b_2 \cos(B_1B_2) + C_5b_3b_4 \cos(B_3B_4) + C_6b_5b_6 \cos(B_5B_6) + C_5C_6b_7b_8 \cos(B_7B_8). \end{aligned}$$

Show that Geometry problem 506 is a special case of the foregoing. Give illustrative example, using triangle or other simple geometric figure, by assuming that some of the sides of the first enneagon are zero.

## SOLUTION BY PROPOSER.

Whenever  $A_1, A_2, \dots$  and  $A_8$  are scalar (or algebraic) quantities, and  $B_1, B_2, \dots$  and  $B_8$  have scalar values corresponding in form to those given in the problem, we have the algebraic identity,

$$\begin{aligned} (C_1^2 + C_5C_2^2 + C_6C_3^2 + C_5C_6C_4^2)(A_1A_2 + C_5A_3A_4 + C_6A_5A_6 + C_5C_6A_7A_8) \\ = B_1B_2 + C_5B_3B_4 + C_6B_5B_6 + C_5C_6B_7B_8. \end{aligned}$$

Inasmuch as all the terms in  $A_1, A_2, \dots, A_8$ , and  $B_1, B_2, \dots, B_8$  in this algebraic identity are of the second degree, a geometric interpretation may be obtained by assuming that  $A_1, A_2, \dots, A_8$  and  $B_1, B_2, \dots, B_8$ , are vectors. This follows immediately from the fact that vector multiplication is commutative in so far as the scalar part of the product is concerned whenever all the vector terms employed are of the second degree. But the scalar part of the vector product  $B_rB_s$  is  $-b_rb_s \cos(B_rB_s)$ . Substituting this scalar part of the vector product in both members of the above algebraic equation and changing signs we obtain at once the equation contained in the problem. If  $A_1 = A_2, A_3 = A_4, A_5 = A_6$ , and  $A_7 = A_8$ , the problem becomes identical with Geometry problem 506.

*Example.* As long as a vector maintains a constant length and direction in space, its origin in space may be altered at will. Hence we need not confine our attention to closed geometric figures in interpreting the given identity. Let  $DEF$  be a given triangle. Bisect  $DE$  in  $G$ . Draw  $GF$  and extend  $EF$  to  $F'$ . Draw  $GH$  intersecting  $DF$  in  $I$  and  $EF'$  in  $J$ . Let  $A_1 = GF, A_2 = GI$ , and  $A_3 = GD$ . Also let  $A_4 = A_5 = A_6 = A_7 = A_8 = 0, C_1 = C_2 = C_3 = C_4 = C_5 = 1$ , and  $C_6 = 0$ . Substituting in the given identity, paying strict attention to the direction of the vectors employed, and dividing by the constant factor, we readily get  $2GF \cos FGI = DF \cos GID + EF \cos EJG$ , a known result.

**2670. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.**

A telegraph wire, weighing one tenth pound per yard, is stretched between poles on level ground, so that the greatest dip of the wire is three feet. Find approximately the distance between the poles when the tension at the lowest point of the wire is 140 pounds.

## SOLUTION BY ELBERT H. CLARKE, Hiram College, Ohio.

It is a well-known property of wires hanging freely from two supports that the tension at